

For Patients

Naloxone works to reverse an overdose of opioids such as fentanyl, heroin, morphine, or oxycodone if given in time.

Responding to an Overdose	Identifying an Overdose
<ol style="list-style-type: none"> 1. Try to wake the person up. 2. Call 911. 3. Administer Naloxone. 4. Repeat a second dose after 2-3 minutes if not breathing. 5. Check for breath. Give CPR or do rescue breathing. 6. Stay with the person. 	<ul style="list-style-type: none"> • Loss of consciousness • Unresponsive to outside stimuli • Awake, but unable to talk • Breathing is slow, erratic, or stopped • Change in skin tone to blue, gray, or ashen • Choking sounds or snore-like gurgling noise

- See full naloxone training video at: <https://www.narcan.com>

For Pharmacists

Naloxone may be available by standing order or dispensed by a pharmacist. Use this prescription if you need a prescription for insurance reasons or other barriers.

Thank you for saving lives.



Dr. Roneet Lev, MD
4077 5th Ave
San Diego, CA 92103
619-203-7000

Patient Name: _____


Date of Birth: _____

Kloxxado Nasal Spray 8 mg/0.1 ml sprayer (Two Pack)

- OR -

Narcan Nasal Spray 4 mg/0.1 ml sprayer (Two Pack)
2 packages

If overdose, call 911. Administer a single spray in the nostril for an opioid overdose. May repeat in 2-3 minutes in the other nostril if patient is not breathing. May substitute as needed for cost or insurance with other formulations of naloxone.

Signature: 
Roneet Lev, MD, FACEP
California license #A48648